

Balancing pastoral and plantation forestry options in New Zealand and the role of agroforestry

Grant B Douglas^A, Andrew J Wall^B, Michael B Dodd^A, Martin F Hawke^C and Ian R McIvor^D

^A AgResearch, Private Bag 11008, Palmerston North, New Zealand, www.agresearch.co.nz

^B AgResearch, Private Bag 50034, Mosgiel, New Zealand, www.agresearch.co.nz

^C Bay of Plenty Farm & Pastoral Research, 23 Glenroy Place, RD 4, Rotorua, New Zealand

^D Plant and Food Research, Private Bag 11600, Palmerston North, New Zealand, www.plantandfood.co.nz

Contact email: grant.douglas@agresearch.co.nz

Abstract. Pastoral agriculture and forestry enterprises are key features of New Zealand's landscape and are very important economically. They are competing landuses, particularly on moderate to steep hill country. Agroforestry involving spaced trees of *Pinus radiata* on pasture was developed in the 1970s to provide dual incomes from livestock enterprises and the later tree crop. In contrast, wide-spaced trees of predominantly *Populus* and *Salix* spp. are planted mainly for erosion control. Characteristics of pastoralism and plantation forestry are reviewed, including trends in conversions between these landuses. Roles, challenges and opportunities with wide-spaced trees are presented, together with current and future research initiatives. Pastoralism and forestry will continue to compete strongly for hill country sites and at present there is an increasing trend of converting previously forested areas to pasture, particularly in the central North Island. Agroforestry involving *Pinus radiata* has virtually ceased because of adverse effects on wood quality, pasture production and animal performance. There are millions of wide-spaced trees of *Populus* and *Salix* spp. on hill country and their planting is expected to continue unabated because they are the most practical and efficient means of enabling pastoralism on erodible slopes and they provide multiple ecosystem services. The species have significant advantages compared to other woody species but many older trees have grown very large because they have received negligible or no silviculture. This is an increasing problem, requiring development and implementation of appropriate management strategies. There is growing interest by landowners in the environmental outcomes of spaced-tree plantings.

Keywords: Soil conservation, Salicaceae, tree-pasture systems, silvopastoralism

Introduction

New Zealand is a hilly and mountainous country with a total land area of 26.8M ha. Over 20% of its land surface exceeds 1,000 m elevation and at least 40% comprises steep, non-arable hill country below 1,000 m elevation (Blaschke *et al.* 1992). At least 1,000 years ago, all of the North Island and most of the South Island were forested (Cameron 1962; White 1999). With habitation by Maori people within the last 800 years, and in particular European settlers within the last 200 years, large tracts of indigenous mixed broadleaved, podocarp and coniferous forests have been cleared for the growing of food crops, energy supply, building materials, and the establishment of a range of pastoral grazing enterprises. Commercial plantation forestry involving exotic tree species commenced in the late 1800s/early 1900s, often in areas that were considered uneconomic or unsuitable for agriculture. Today, low to highly productive grassland covers about 11.0M ha (41%) of the country's land area (Beef + Lamb New Zealand 2012) and plantation forestry comprises 1.75M ha (7%) (Hedley *et al.* 2009, Anon 2012). Both sectors contribute significantly to New Zealand's gross domestic product (Ministry of Agriculture and Forestry 2011).

Plantation forests are distributed widely throughout New Zealand but most of the area (70.3%) occurs in the North Island, almost half of which is in the central regions (New Zealand Forest Owners Association 2011). Most of the forests are grown on moderate to steep hill country (Nagashima *et al.* 2001; Quinn *et al.* 2007; Raymond 2012) with slope angles often exceeding 20°. These areas are unsuitable for arable use and achievement of long-term sustainable productivity can be compromised because of factors such as moderate to severe mass movement erosion, shallow soils with increasing steepness, low soil organic matter status and poor water holding capacity, soil nutrient deficiencies and scrub weed reversion (Gillingham *et al.* 1998; Matthews *et al.* 1999; Dodd *et al.* 2008; De Rose 2012).

The New Zealand landscape has been classified on the basis of its land use capability (LUC) into classes 1 to 8 according to its capacity for long-term sustained production (NWASCO 1979, Lynn *et al.* 2009). As class number increases, there are increasing limitations to use of the land and a corresponding decrease in its versatility of use. Limitations may include unfavourable climate, erosion risk, shallow soils, low soil water holding capacity, stoniness, and excessive wetness. About 7.5M ha (28%) of New Zealand

and has been mapped as LUC Class 6, most of which is stable productive hill country with moderate erosion potential. LUC Class 7 comprises 5.7M ha (21%) of the country and has limitations similar to those for Class 6 land, but they are more severe. LUC Class 6 land is suitable for both pastoralism and forestry whereas Class 7 land is more suitable for forestry in many instances (Lynn *et al.* 2009). However with the implementation of significant soil conservation measures, Class 7 land can be suitable for pastoralism. Most plantation forestry occurs on LUC Classes 6 and 7, rather than the more versatile Classes 1-5. LUC Class 8 land is unsuitable for pastoralism or forestry enterprises because of very severe to extreme limitations, often extreme actual or potential erosion (Lynn *et al.* 2009), and is used mainly for conservation of flora and fauna, erosion control and water management.

Pastoral enterprises and plantation forestry have been competing landuses on moderate to steep hill country (LUC Classes 6 and 7) for many decades. The temporal and spatial mosaic of these landuses on these landscapes has been governed by numerous factors including capital outlay and economic returns over various timeframes, environmental effects, political and regulatory initiatives, and social and cultural considerations (Ross *et al.* 2002, Hawke 2004, Douglas *et al.* 2006a, Elliott and Hawke 2007, Lee 2007, Hedley *et al.* 2009). Competition between these landuses occurs mostly at the individual farm level but can occur at other scales (*e.g.* district) and have quite far-reaching effects such as on water quality at the catchment scale, and the well-being of local communities. The development of agroforestry in the 1970s aimed for both livestock production from pastures, and timber production from wide-spaced trees, usually *Pinus radiata*, on the same land unit (Hawke 1991, Knowles 1991, Hawke and Wedderburn 1994). The term agroforestry may include additional systems such as forests/woodlots and shelterbelts, but is confined here to wide-spaced tree plantings. Other tree-pasture systems, usually based on *Populus* or *Salix* spp., are established on large tracts of erodible hill country to enable the continuation of pastoralism (Wilkinson 1999; Benavides *et al.* 2009). However in these systems, tree spacing is often highly variable and targeted to areas deemed of greatest erosion risk. Furthermore, the trees are planted mainly for soil conservation rather than timber production, and consequently they have usually received negligible or no silviculture (Cameron 2003).

The objectives of this paper are to outline key features of pastoralism and forestry in New Zealand hill country, trends in conversions between these landuses, background to agroforestry and the extent of its current use, and the role of wide-spaced trees. Finally, we describe current and planned research initiatives with tree-pasture systems.

Pastoralism

Developed livestock grazing systems in New Zealand hill country are based on year-round grazing of predominantly temperate, introduced grasses, legumes and herbs (Kemp *et al.* 1999, Charlton and Stewart 2006). *Lolium perenne* and *Trifolium repens* are components of nearly all seed mixtures but a range of other species may be included (Scott *et al.* 1985; Kemp *et al.* 1999). Hill country pastures are

farmed in development states ranging from low (unimproved) to high depending on aspects including fertiliser strategies, extent of species introduction and general over-sowing practices, stocking rate and pasture utilisation, and effectiveness of weed and animal pest control programmes. Annual utilisable pasture production in North and South Island hill country is mostly in the range 5-10 t dry matter (DM)/ha, although 20-25 t DM/ha/yr can be achieved on livestock camps in moist areas (Chapman and Macfarlane 1985).

Livestock on hill country farms are predominantly sheep and beef cattle, with farms in the South Island carrying a greater ratio of sheep:cattle compared with North Island farms (Matthews *et al.* 1999). Deer are also farmed in some hill country enterprises for venison and antlers/velvet. Environmental challenges and consequences of pastoral farming of hill country include: loss of nutrients originating from grazing animals and fertiliser applications through overland flow and leaching; contamination of water bodies with sediment and nutrients; contributions to greenhouse gas (GHG) emissions; changes to soil invertebrate populations; and potential damage to soil structure and other soil physical characteristics (Betteridge *et al.* 1999, Russell *et al.* 2001, Schon *et al.* 2011, Mackay *et al.* 2012b). There can also be total loss of soil profile following extreme erosion events. Pastoral hill country is up to 20-times more susceptible to mass movement erosion than similar areas with woody vegetation (Hicks 1991, Marden and Rowan 1993, Dymond *et al.* 2006), depending mainly on lithology, slope, and rainfall.

Plantation forestry

The predominant tree species grown in New Zealand is *Pinus radiata*, comprising approximately 90% of the area of the national estate (New Zealand Forest Owners Association 2011). The species is hardy, fast-growing, adapted to a variety of environmental conditions, productivity often exceeds 25 m³/ha/yr in many parts of the country and trees are ready to harvest in 25-30 years (Kimberley *et al.* 2005; Kirschbaum *et al.* 2012; Woollons and Manley 2012). Other species grown include *Pseudotsuga menziesii*, *Cupressus macrocarpa*, *Acacia melanoxylon* and *Eucalyptus* spp.

In addition to their production of wood for a range of products and purposes, hill slopes with established forests have lower susceptibility to erosion than with any other ground cover, and this appears largely independent of species (Phillips *et al.* 1990; Brown 1991; Cairns *et al.* 2001; Dymond *et al.* 2006). The presence of forests enhances other ecosystem services such as flood mitigation, carbon storage in wood and soil, filtering of nutrients, and even cultural services such as recreation. Erosion management in tree plantations is critical at establishment and harvesting, where vegetation removal and mechanical earthworks temporarily increase soil movement and the risk of significant soil loss (Marden *et al.* 2005, 2006). Extensive guidelines (Vaughan 1984; Spiers 1987) and a code of practice (New Zealand Forest Owners Association 2007) have been developed to guide operational planning and maintain soil and water values through mitigation of erosion and sedimentation.

The role of exotic forests in sequestering carbon to reduce atmospheric GHG concentrations has received increasing attention, both in terms of environmental management and revenue generation (Hollinger *et al.* 1993; Robertson *et al.* 2004; Adams and Turner 2012). In a recent study of *Pinus radiata* at 140 sites, it was concluded that mainly untended trees could be grown on a rotation length of 60+ years to enable a longer period of carbon sequestration than from contemporary rotation lengths (Woollons and Manley 2012). Such an approach requires appraisal in the context of factors including carbon market price, tree stand area, and projected returns as trees age.

Conversions between pastoralism and forestry

During the 1990s, 20,000-90,000 ha of pastoral land was converted to plantation forestry annually, and concomitant research determined the impacts of this land use change, predominantly with respect to soil chemistry (Alfredsson *et al.* 1998; Ross *et al.* 1999; Parfitt and Ross 2011), but also other soil attributes including soil bacteria, bulk density and hydrophobicity (Singh *et al.* 2007; Paripovic 2011). Since the mid-1990s, there was a significant reduction in new plantings mainly because of reduced numbers of new conversions from pasture, with the lowest area of 1,900 ha planted in 2008 (Ministry for Primary Industries 2012). In the mid-2000s, there was also an increase in the practice of not replanting forests after harvest and in a few cases, forests were converted to pasture before they reached maturity (Ministry for the Environment 2007). Recently, the area of new plantings has increased, partly because of grants for afforestation as part of the government's Emissions Trading Scheme (<http://www.climate-change.govt.nz/emissions-trading-scheme/about/basics.html>), with an estimated 12,000 ha of new plantings in 2011 and a provisional estimate of 18,500 ha in 2012 (Ministry for Primary Industries 2012).

In the last decade, deforestation of exotic forests and conversion to pastoralism has been an increasing trend, mainly in the central North Island for dairy production (Hawke 2004; Elliott and Hawke 2007; Hedley *et al.* 2009; Paripovic 2011). A perennial pasture was established successfully on an ex-forest site with addition of lime (4 t/ha) and phosphorus (40 kg/ha) to provide an appropriate pH and Olsen P status for grass and legume growth (Hawke 2004). In the region, Brodnax (2007) estimated that the total conversion may be as high as 60,000 ha, and will have a number of environmental consequences for local and regional water systems, principally because of the significant dung and urine excretions from the introduced livestock. Hydrology of the catchment will also be modified because of increased runoff (lack of flood mitigation service).

Agroforestry

Agroforestry systems in New Zealand have been evaluated since the 1970s with respect to the tree component, pasture understorey, soils, animal performance, and their interactions, continuing through the 1990s (Hawke 1991; Knowles 1991; Hawke *et al.* 1993; Hawke and Wedderburn 1994; Hawke 1997; Knowles *et al.* 1999; Maclaren and Knowles 1999a; Perrott *et al.* 1999) and into this century (Amatya *et al.* 2002; Peri *et al.* 2002; Chang and Mead 2003;

Benavides *et al.* 2009). The Tikitere project in the central North Island was a major source of knowledge on the impacts of tree density (tree stocking rate) (50, 100, 200 and 400 stems per hectare (sph)) on the system components. For example, trees at lower densities had considerably larger diameter growth, greater stem and branch size, and were shorter than those at higher densities (Hawke 2011). When the trees were harvested at 25 years, total volume ranged from 213 m³/ha at 50 sph to 935 m³/ha at 400 sph. Canopy closure occurred earlier at high than low tree densities with high densities causing the greatest reduction in pasture understorey production and change in botanical composition. Hawke (2011) reported that increasing tree density and age resulted in reduced animal performance. Despite much experience with agroforestry in New Zealand, there has been negligible recent reporting of economic aspects (Maclaren and Knowles 1999b, Hawke 2011).

Current interest in agroforestry has waned considerably and we are unaware of any new blocks being established on farms. A recent harvest of *Pinus radiata* at 155 sph concluded that there was a lack of volume of logs from such a system compared with trees grown in plantation at 320 sph and that "you cannot, it seems, have it both ways" (Jones and Cullen 2008). This was reinforced by the findings from the Tikitere project where it was concluded that agroforestry adversely affected wood quality, pasture production and animal performance (Hawke 2011). There does not appear to be a future for agroforestry involving *Pinus radiata* but there may be opportunities with high-value timber species such as some belonging to the genera *Cupressus*, *Eucalyptus*, *Quercus* and *Sequoia*, and species that are deciduous, with less impact on pasture understorey growth.

Wide-spaced trees to support pastoralism on hill country

There are millions of wide-spaced plantings of *Populus* and *Salix* spp. on erodible pastoral hill country, particularly where erosion arises from surplus soil water (Thompson and Luckman 1993; Wilkinson 1999). Key advantages of these species include: an ability to establish in the presence of grazing livestock (sheep and sometimes young cattle) using vegetative cuttings with plastic sleeve protectors; deciduous growth habit; extensive and rapidly developing root systems (McIvor *et al.* 2008; McIvor *et al.* 2009; Douglas *et al.* 2010); and effectiveness in reducing a range of erosion processes (Thompson and Luckman 1993; Douglas *et al.* 2013). They also provide shade and shelter for grazing livestock, valuable supplementary fodder during times of soil water deficit, vista enhancement, potential for gaining carbon credits, flood mitigation by intercepting rainfall and increasing drainage, and biodiversity (McGregor *et al.* 1999; Kemp *et al.* 2001; Kemp *et al.* 2003; Betteridge *et al.* 2012; McIvor and Douglas 2012). Numerous clones are available, varying in attributes including growth form, leaf retention, tolerance to wind, and pest and disease resistance (National Poplar and Willow Users Group 2007). On seasonally drought-prone sites, such as those on north-facing upper slopes, where *Populus* spp. and *Salix* spp. may fail to establish or survive, *Acacia* spp. and *Eucalyptus* spp. are options (Sheppard and Bulloch 1986; Bulloch 1991; Millner and Kemp 2012).

Trees on slopes should be spaced to provide effective soil stabilisation and enable adequate light transmission for understorey pasture growth. Interlocking of roots from neighbouring trees, analogous to overlaying a steel reinforcing mesh across a planted slope, is critical for enhancing soil strength and resistance to failure. Wilkinson (1999) advocated hillslope plantings of *Populus* and *Salix* at densities of 25 to 100 sph, equivalent to spacings of 20 m and 10 m, respectively. On the East Coast of the North Island, individual trees of *Populus x euramericana* aged 14–17 years and spaced 20 m apart, on average, saved 8.4 m² of ground from shallow landsliding following a severe storm (Hawley and Dymond 1988). Across all trees in their study, the area of landslide scar was reduced by 13.8% compared with similar sites without trees, and modelling suggested that at a tree spacing of 10 m, the reduction would have been at least 70%. In the lower North Island on slopes averaging 27°, established trees at 32–65 sph (18–12 m spacing), mostly *Populus* spp., reduced shallow landslide occurrence by 95% compared to paired pasture control sites (Douglas et al. 2013). Excavations of entire root systems indicated that trees need to be older than 5 years to make worthwhile contributions to slope reinforcement through lateral and vertical root development (McIvor et al. 2008, McIvor et al. 2009). It is recommended to plant *Populus* spp. at 8–15 m centres depending on the severity of erosion (National Poplar and Willow Users Group 2007), with thinning to achieve wider spacing possible as trees mature.

The impact of wide-spaced trees on understorey pasture yield and quality has received considerable attention for more than a decade, mainly in relation to *Populus* spp. (Guevara-Escobar et al. 1997; Douglas et al. 2001; Douglas et al. 2006b; Wall 2006; Wall et al. 2006; Guevara-Escobar et al. 2007; Benavides et al. 2009; Wall et al. 2010), but also beneath and beyond trees of *Acacia*, *Alnus*, *Eucalyptus* and *Salix* spp. (Gilchrist et al. 1993; Power et al. 1999, 2001; Devkota et al. 2009). Under mature, unpruned *Populus deltoides* (mean diameter at breast height (DBH) = 70 cm) at a spacing averaging 16.4 m (37.2 sph), providing essentially 100% canopy closure during foliage, annual pasture accumulation was reduced by 40% compared with nearby areas without trees (Guevara-Escobar et al. 2007). Accumulation beneath younger trees was similar beneath and beyond 5-year-old trees (Guevara-Escobar et al. 2007), 7 or 11% less beneath trees aged 7 years than beyond (McIvor and Douglas 2012), and 23% less beneath trees aged 8–11 years than beyond (Douglas et al. 2006b). Modelling indicated that pasture yield during summer beneath *Populus x euramericana* trees aged 16–19 years was 12.5–21% less than on areas without trees (Gilchrist et al. 1993). There has been a burgeoning development and understanding of canopy closure models and impacts (McElwee and Knowles 2000; Guevara-Escobar et al. 2005; Wall et al. 2006; Wall et al. 2010), reflecting the major influence of radiation interception on understorey responses (Dodd et al. 2005). These will prove useful for evaluating tree spacing scenarios and their likely implications.

Populus and *Salix* species are important and widely used sources of timber and other products internationally (Turley et al. 2006; Spinelli et al. 2011) but the potential of wide-spaced plantings in New Zealand has not been realised, despite various attempts (McGregor et al. 1999). The

general lack of silviculture in any form, and no harvesting, has resulted in many of the older plantings developing into very large trees e.g. DBH > 60 cm, which significantly shade understorey pasture and are a liability to farm infrastructure, livestock and humans through potential limb breakage and toppling. Few options exist for dealing with large trees which has deterred some current farmers from commencing or continuing planting programmes. Apart from removal by professional logging contractors, large trees can be poisoned by injecting the trunk with herbicide, which results in death and decay but without certainty of the timing of these events. Glyphosate and metsulfuron-methyl were more effective in killing old *Populus* trees (DBH = 70–101 cm) when injected between October and February during their active growth, than when injected in April when growth had slowed (National Poplar and Willow Users Group 2007), though all trees died eventually.

Pollarding, involving complete removal of the tree canopy, particularly at young and intermediate tree ages, is practised by some farmers for limiting tree size and reducing shading of pasture (National Poplar and Willow Users Group 2007). Unpublished results from a study determining the effect of pollarding trees of *Populus deltoides* x *P. nigra* clone 'Veronese' (mean DBH = 21 cm) found that 4 years after pollarding, the increase in DBH of pollarded trees (3.3 cm) was 66% less than for un-pollarded trees (9.8 cm), and coarse root (> 2 mm diameter) mass and length were about 60% less for pollarded compared with un-pollarded trees. These findings have important implications for slope stabilisation because they suggest that pollarded trees need to be spaced closer together than un-pollarded trees to achieve similar levels of effectiveness.

There have been very few economic analyses of wide-spaced trees in pastoral systems (Parminter et al. 2001; Orsborn et al. 2003). A cost-benefit analysis of planting *Populus* spp. on erosion-prone hill country under three scenarios, varying mainly in erosion severity, gross margin per stock unit, and fodder value, found that the on-site profitability of planting for erosion control alone was marginal at best (Parminter et al. 2001). It was suggested that factors such as likely reduced lamb losses, reduced off-site effects, vista enhancement and reduced insurance payouts from storm damage, were added incentives for planting programmes. On a hill country farm, modelling predicted a financial advantage from feeding ewes *Populus* and *Salix* foliage as a supplement to pasture during dry conditions in summer/early autumn (Orsborn et al. 2003).

Current and new research initiatives for wide-spaced trees

Following an extensive literature review, ten aspects were deemed important for future understanding and knowledge of the complexities and functions of tree-pasture systems (Benavides et al. 2009). These were effectiveness of trees for controlling erosion, their root distribution, controlling tree size, modelling system impacts on erosion processes, effect of canopy shape on pasture understorey growth and links with root distribution and water use, alternative tree species, carbon sequestration, spaced-tree effects on water, nutrient and sediment patterns, animal welfare, and eco-

nomic analyses.

Some of these aspects have been addressed recently including the effectiveness of wide-spaced trees in reducing the occurrence of shallow landslides on steep slopes (Douglas *et al.* 2013), increased understanding of root distribution (McIvor *et al.* 2009; Douglas *et al.* 2010), and the effect of spaced trees on grazing behaviour and physiology of beef cattle (Betteridge *et al.* 2012). New or on-going initiatives include those determining the impact of pollarding trees of *Populus* and *Salix* spp. on understorey pasture production, water use patterns, and root development (McIvor *et al.* 2011), linking vegetation management on-farm, including spaced-tree conservation plantings, with reducing the risk of erosion and sediment loss to waterways (Mackay *et al.* 2012a), and the effect of *Populus* and *Alnus* spp. at different densities (60-1,200 sph) on soil carbon stocks to 1 m depth. Environmental outcomes of tree-pasture systems on hill country will be of increasing interest.

Conclusion

Strong competition between landuses of forestry and pastoralism on moderate to steep hill country is on-going and the balance between them is governed by a wide range of socio-economic and other factors. The momentum gained in the 1970s-1990s in blending the two landuses on the same land unit in the form of agroforestry with *Pinus radiata*, has waned considerably, to the extent that few, if any, new examples exist on farms. Other tree species may be considered in future agroforestry initiatives. There has and continues to be active planting of wide-spaced trees of *Populus* and *Salix* spp., primarily for erosion control, but also for a range of other functions. Research is addressing some key aspects of these ecosystems.

Acknowledgements

This review was funded by the Sustainable Landuse Research Initiative programme, funded by Ministry of Business, Innovation and Employment (core-funding to the New Zealand Institute for Plant and Food Research Limited).

References

- Adams T, Turner JA (2012). An investigation into the effects of an emissions trading scheme on forest management and land use in New Zealand. *Forest Policy and Economics* **15**, 78-90.
- Alfredsson H, Condon LM, Clarholm M, Davis MR (1998). Changes in soil acidity and organic matter following the establishment of conifers on former grassland in New Zealand. *Forest Ecology and Management* **112**, 245-252.
- Amatya G, Chang SX, Beare MH, Mead DJ (2002). Soil properties under a *Pinus radiata* - ryegrass silvopastoral system in New Zealand. Part II. C and N of soil microbial biomass, and soil N dynamics. *Agroforestry Systems* **54**, 149-160.
- Anon (2012). 'Forestry and the New Zealand Economy.' Available at www.mpi.govt.nz/forestry [Accessed 19 October, 2012].
- Beef + Lamb New Zealand (2012). Compendium of New Zealand farm facts. Beef + Lamb New Zealand Ltd. Economic Service, Wellington. No. P12009.
- Benavides R, Douglas GB, Osoro K (2009). Silvopastoralism in New Zealand: review of effects of evergreen and deciduous trees on pasture dynamics. *Agroforestry Systems* **76**, 327-350.
- Betteridge K, Costall D, Martin S, Reidy B, Stead A, Millner I (2012). Impact of shade trees on Angus cow behaviour and physiology in summer dry hill country: grazing activity, skin temperature and nutrient transfer issues. In 'Advanced Nutrient Management: Gains from the Past - Goals for the Future.' (Eds LD Currie, CL Christensen.) pp. 10. (Fertilizer and Lime Research Centre, Massey University: Palmerston North)
- Betteridge K, Mackay AD, Shepherd TG, Barker DJ, Budding PJ, Devantier BP, Costall DA (1999). Effect of cattle and sheep treading on surface configuration of a sedimentary hill soil. *Australian Journal of Soil Research* **37**, 743-760.
- Blaschke PM, Trustrum NA, DeRose RC (1992). Ecosystem processes and sustainable land use in New Zealand steepplands *Agriculture, Ecosystems & Environment* **41**, 153-178
- Brodnax R (2007). The next chapter in regional policy development to manage adverse effects of land use intensification. In 'Designing Sustainable Farms: Critical Aspects of Soil and Water Management.' (Eds LD Currie, LJ Yates.) pp. 41-49. (Fertilizer and Lime Research Centre, Massey University: Palmerston North)
- Brown WJ (1991). Landslide control on North Island, New Zealand. *Geographical Review* **81**, 457-472.
- Bulloch BT (1991). *Eucalyptus* species selection for soil conservation in seasonally dry hill country - twelfth year assessment. *New Zealand Journal of Forestry Science* **21**, 10-31.
- Cairns I, Handyside B, Harris M, Lambrechtsen N, Ngapo N (Eds DL Hicks, T Anthony (2001). 'Soil conservation technical handbook.' (New Zealand Association of Resource Management: Wellington)
- Cameron DJ (2003). Trees and pasture - An overview. In 'Using Trees on Farms.' (Ed. JFL Charlton.) pp. 5-6. (New Zealand Grassland Association: Wellington)
- Cameron RJ (1962). The exploitation of the indigenous forests. *Proceedings of the New Zealand Ecological Society* **9**, 50-56.
- Chang SX, Mead DJ (2003). Growth of radiata pine (*Pinus radiata* D. Don) as influenced by understory species in a silvopastoral system in New Zealand. *Agroforestry Systems* **59**, 43-51.
- Chapman DF, Macfarlane MJ (1985). Pasture growth limitations in hill country and choice of species. In 'Using Herbage Cultivars.' (Eds RE Burgess, JL Brock.) pp. 25-29. (Grassland Research and Practice Series No. 3, New Zealand Grassland Association: Palmerston North)
- Charlton D, Stewart A (2006). 'Pasture and forage plants for New Zealand.' (New Zealand Grassland Association/ New Zealand Grassland Trust: Wellington)
- De Rose RC (2012). Slope control on the frequency distribution of shallow landslides and associated soil properties, North Island, New Zealand. *Earth Surface Processes and Landforms* in press.
- Devkota NR, Kemp PD, Hodgson J, Valentine I, Jaya IK (2009). Relationship between tree canopy height and the production of pasture species in a silvopastoral system based on alder trees. *Agroforestry Systems* **76**, 363-374.
- Dodd MB, McGowan AW, Power IL, Thorrold BS (2005). Effects of variation in shade level, shade duration and light quality on perennial pastures. *New Zealand Journal of Agricultural Research* **48**, 531-543.
- Dodd MB, Wedderburn ME, Parminter TG, Thorrold BS, Quinn JM (2008). Transformation toward agricultural sustainability in New Zealand hill country pastoral landscapes. *Agricultural Systems* **98**, 95-107.
- Douglas G, Mackay A, Dodd M, Rhodes T, Budding P (2006a). Land-use intentions and changes in shrubland cover on farms in the Wanganui region since the 1960's. AgResearch Ltd, Palmerston North.
- Douglas GB, McIvor IR, Manderson AK, Koolaard JP, Todd M,

- Braaksma S, Gray RAJ (2013). Reducing shallow landslide occurrence in pastoral hill country using wide-spaced trees. *Land Degradation and Development* **24**, 103-114.
- Douglas GB, McIvor IR, Potter JF, Foote LG (2010). Root distribution of poplar at varying densities on pastoral hill country. *Plant and Soil* **333**, 147-161.
- Douglas GB, Walcroft AS, Hurst SE, Potter JF, Foote AG, Fung LE, Edwards WRN, van den Dijssel C (2006b). Interactions between widely spaced young poplars (*Populus* spp.) and introduced pasture mixtures. *Agroforestry Systems* **66**, 165-178. doi: [10.1007/s10457-005-6641-5](https://doi.org/10.1007/s10457-005-6641-5)
- Douglas GB, Walcroft AS, Wills BJ, Hurst SE, Foote AG, Trainor KD, Fung LE (2001). Resident pasture growth and the micro-environment beneath young, wide-spaced poplars in New Zealand. *Proceedings of the New Zealand Grassland Association* **62**, 131-138.
- Dymond JR, Ausseil A-G, Shepherd JD, Buettner L (2006). Validation of a region-wide model of landslide susceptibility in the Manawatu-Wanganui region of New Zealand. *Geomorphology* **74**, 70-79.
- Elliott I, Hawke MF (2007). Best management practices for forestry to farming conversions. *Proceedings of the New Zealand Grassland Association* **69**, 79-86.
- Gilchrist AN, Hall JR, Foote AG, Bulloch BT (1993). Pasture growth around broadleaved trees planted for grassland stability. *Proceedings of XVII International Grassland Congress Vol. III, Rockhampton, Australia 2062-2063*.
- Gillingham AG, Gray MH, Smith DR (1998). Pasture responses to phosphorus and nitrogen fertilisers on dry hill country. *Proceedings of the New Zealand Grassland Association* **60**, 135-140.
- Guevara-Escobar A, Kemp PD, Hodgson J, Mackay AD, Edwards WRN (1997). Case study of a mature *Populus deltoides*-pasture system in a hill environment. *Proceedings of the New Zealand Grassland Association* **59**, 179-185.
- Guevara-Escobar A, Kemp PD, Mackay AD, Hodgson J (2007). Pasture production and composition under poplar in a hill environment in New Zealand. *Agroforestry Systems* **69**, 199-213.
- Guevara-Escobar A, Tellez J, Gonzalez-Sosa E (2005). Use of digital photography for analysis of canopy closure. *Agroforestry Systems* **65**, 175-185.
- Hawke M (2011). Agroforestry - the Tikitere experience. *New Zealand Tree Grower* **32**, 23-26.
- Hawke MF (1991). Pasture production and animal performance under pine agroforestry in New Zealand. *Forest Ecology and Management* **45**, 109-118.
- Hawke MF (1997). The changing focus of pastoral agroforestry in temperate zones. *XVIII International Grassland Congress, June 8-19, 1997 Winnipeg, Manitoba, Canada, Session 6: 121-128*.
- Hawke MF (2004) Conversion of forestry land back to productive pasture. *Proceedings of the New Zealand Grassland Association* **66**, 157-162.
- Hawke MF, Rattray PV, Percival NS (1993). Liveweight changes of sheep grazing a range of herbage allowances under *Pinus radiata* agroforestry regimes. *Agroforestry Systems* **23**, 11-21.
- Hawke MF, Wedderburn ME (1994). Microclimate changes under *Pinus radiata* agroforestry regimes in New Zealand. *Agricultural and Forest Meteorology* **71**, 133-145.
- Hawley JG, Dymond JR (1988). How much do trees reduce landsliding? *Journal of Soil and Water Conservation* **43**, 495-498.
- Hedley CB, Kusumo BH, Hedley MJ, Tuohy MP, Hawke M (2009). Soil C and N sequestration and fertility development under land recently converted from plantation forest to pastoral farming. *New Zealand Journal of Agricultural Research* **52**, 443-453.
- Hicks DL (1991). Erosion under pasture, pine plantations, scrub and indigenous forest: a comparison from Cyclone Bola. *New Zealand Forestry* **36**, 21-22.
- Hollinger DY, Maclaren JP, Beets PN, Turland J (1993). Carbon sequestration by New Zealand's plantation forests. *New Zealand Journal of Forestry Science* **23**, 194-208.
- Jones K, Cullen N (2008). Wide planting does not work. *New Zealand Tree Grower* **29**, 4.
- Kemp PD, Barry TN, Douglas GB (2003). Edible forage yield and nutritive value of poplar and willow. In 'Using Trees on Farms.' (Ed. JFL Charlton.) pp. 53-63. (New Zealand Grassland Association: Wellington, New Zealand)
- Kemp PD, Mackay AD, Matheson LA, Timmins ME (2001). The forage value of poplars and willow. *Proceedings of the New Zealand Grassland Association* **63**, 115-119.
- Kemp PD, Matthew C, Lucas RJ (1999). Pasture species and cultivars. In 'New Zealand pasture and crop science.' (Eds J White, J Hodgson.) pp. 83-99. (Oxford University Press: Auckland)
- Kimberley M, West G, Dean M, Knowles L (2005). The 300 index - A volume productivity index for radiata pine. *New Zealand Journal of Forestry* **50**, 13-18.
- Kirschbaum MUF, Watt MS, Tait A, Ausseil A-GE (2012). Future wood productivity of *Pinus radiata* in New Zealand under expected climatic changes. *Global Change Biology* **18**, 1342-1356. doi: [10.1111/j.1365-2486.2011.02625.x](https://doi.org/10.1111/j.1365-2486.2011.02625.x)
- Knowles RL (1991). New Zealand experience with silvopastoral systems: A review. *Forest Ecology and Management* **45**, 251-267. doi: [10.1016/0378-1127\(91\)90221-g](https://doi.org/10.1016/0378-1127(91)90221-g)
- Knowles RL, Horvath GC, Carter MA, Hawke MF (1999). Developing a canopy closure model to predict overstorey/understorey relationships in *Pinus radiata* silvopastoral systems. *Agroforestry Systems* **43**, 109-119.
- Lee B (2007). Sustainable land use: a farm forester's perspective. *Proceedings of the New Zealand Grassland Association* **69**, 107-109.
- Lynn IH, Manderson AK, Page MJ, Harmsworth GR, Eyles GO, Douglas GB, Mackay AD, Newsome PJF (2009). 'Land Use Capability Survey Handbook - a New Zealand handbook for the classification of land.' (AgResearch Ltd, Hamilton, Landcare Research New Zealand Ltd, Lincoln, Institute of Geological and Nuclear Sciences Ltd, Lower Hutt):
- Mackay A, Douglas G, Wheeler D, Power I (2012a). Establishing a common framework for Regional Councils to assess, quantify and report on the effectiveness of soil conservation works on farm erosion risk and potential sediment loss. Report prepared for Horizons Regional Council, Envirolink Medium Advice Grant Project HZLC87. AgResearch Ltd, Palmerston North.
- Mackay AD, Rhodes AP, Power I, Wedderburn ME (2012b). Has the eco-efficiency of sheep and beef farms changed in the last 20 years? *Proceedings of the New Zealand Grassland Association* **74**, 11-16.
- Maclaren JP, Knowles RL (1999a). Economics of final crop stocking at the Tikitere Agroforestry Trial. Part 1: volume and quality comparisons. *New Zealand Journal of Forestry Science* **29**, 165-174.
- Maclaren JP, Knowles RL (1999b). Economics of final crop stocking at the Tikitere Agroforestry Trial. Part 2: Economic comparisons. *New Zealand Journal of Forestry Science* **29**, 175-187.
- Marden M, Arnold G, Gomez B, Rowan D (2005). Pre- and post-reforestation gully development in Mangatu Forest, East Coast, North Island, New Zealand. *River Research and Applications* **21**, 757-771.
- Marden M, Rowan D (1993). Protective value of vegetation on tertiary terrain before and during Cyclone Bola, East Coast, North Island, New Zealand. *New Zealand Journal of Forestry Science* **23**, 255-263.

- Marden M, Rowan D, Phillips C (2006). Sediment sources and delivery following plantation harvesting in a weathered volcanic terrain, Coromandel Peninsula, North Island, New Zealand. *Australian Journal of Soil Research* **44**, 219-232. doi:10.1071/SR05092
- Matthews PNP, Hodgson J, White JGH (1999). Livestock farming systems in New Zealand. In 'New Zealand Pasture and Crop Science.' (Eds J White, J Hodgson.) pp. 133-151. (Oxford University Press: Auckland)
- McElwee HF, Knowles RL (2000). Estimating canopy closure and understorey pasture production in New Zealand-grown poplar plantations. *New Zealand Journal of Forestry Science* **30**, 422-435.
- McGregor E, MacKay AD, Dodd MB, Kemp PD (1999). Silvopastoralism using tended poplars on New Zealand hill country: The opportunities. *Proceedings of the New Zealand Grassland Association* **61**, 85-89.
- McIvor I, Gosling M, Mason K, Gautellier-Vizioz L (2011). Effect of pollarding for stock fodder on root dynamics of conservation willow trees in New Zealand. In 'CRA - Open Conference System, Fifth International Poplar Symposium (IPS V). Available at <http://ocs.entecra.it/index.php/IPS/5/paper/view/139>
- McIvor IR, Douglas GB (2012). Poplars and willows in hill country - stabilising soils and storing carbon. In 'Advanced Nutrient Management: Gains from the Past - Goals for the Future.' (Eds LD Currie, CL Christensen.) pp. 1-11. (Fertilizer and Lime Research Centre, Massey University: Palmerston North)
- McIvor IR, Douglas GB, Benavides R (2009). Coarse root growth of Veronese poplar trees varies with position on an erodible slope in New Zealand. *Agroforestry Systems* **76**, 251-264. doi:10.1007/s10457-009-9209-y
- McIvor IR, Douglas GB, Hurst SE, Hussain Z, Foote AG (2008). Structural root growth of young Veronese poplars on erodible slopes in the southern North Island, New Zealand. *Agroforestry Systems* **72**, 75-86. doi:10.1007/s10457-007-9090-5
- Millner JP, Kemp PD (2012). Seasonal growth of *Eucalyptus* species in New Zealand hill country. *New Forests* **43**, 31-44. doi:10.1007/s11056-011-9264-x
- Ministry for Primary Industries (2012). National Exotic Forest Description as at 1 April 2012.
- Ministry for the Environment (2007). Land. In 'Environment New Zealand 2007.' pp. 210-257. (Ministry for the Environment: Wellington)
- Ministry of Agriculture and Forestry (2011). Situation and Outlook for New Zealand Agriculture and Forestry. Ministry of Agriculture and Forestry, Wellington.
- Nagashima K, Sands R, Whyte AGD, Bilek EM, Nakagoshi N (2001). Forestry expansion and land-use patterns in the Nelson Region, New Zealand. *Landscape Ecology* **16**, 719-729. doi:10.1023/a:1014481114671
- National Poplar and Willow Users Group (2007). 'Growing poplar and willow trees on farms. Guidelines for establishing and managing poplar and willow trees on farms.' (National Poplar and Willow Users Group: Palmerston North)
- New Zealand Forest Owners Association (2007). 'New Zealand environmental code of practice for plantation forestry.' (New Zealand Forest Owners Association: Wellington)
- New Zealand Forest Owners Association (2011). 'New Zealand plantation forest industry facts and figures 2010/11.' (New Zealand Forest Owners Association: Wellington)
- NWASCO (1979). 'Our land resources. Bulletin to accompany New Zealand Land Resource Inventory Worksheets.' (National Water and Soil Conservation Organisation, Ministry of Works and Development: Wellington)
- Orsborn S, Gawith PW, Cameron DJ (2003). Cost-benefits of supplementing ewes with willow and poplar foliage on a model hill country farm in Wairarapa. In 'Using Trees on Farms.' (Ed. JFL Charlton.) pp. 35-40. (Grassland Research and Practice Series No. 10. New Zealand Grassland Association: Wellington, New Zealand)
- Parfitt RL, Ross DJ (2011). Long-term effects of afforestation with *Pinus radiata* on soil carbon, nitrogen, and pH: a case study. *Soil Research* **49**, 494-503. doi: <http://dx.doi.org/10.1071/SR11106>
- Paripovic D (2011). Impacts of conversion from forestry to pasture on soil physical properties of Vitrands (Pumice Soils) in the Central North Island, New Zealand. MSc thesis, University of Waikato.
- Parminter I, Dodd MB, Mackay AD (2001). Economic analysis of poplar planting on steep hill country. *Proceedings of the New Zealand Grassland Association* **63**, 127-130.
- Peri PL, Mason EG, Pollock KM, Varella AC, Mead DJ (2002). Early growth and quality of radiata pine in a silvopastoral system in New Zealand. *Agroforestry Systems* **55**, 207-219.
- Perrott KW, Ghani A, O'Connor MB, Waller JE, Hawke MF (1999). Tree stocking effects on soil chemical and microbial properties at the Tikitere agroforestry research area. *New Zealand Journal of Forestry Science* **29**, 116-130.
- Phillips CJ, Marden M, Pearce AJ (1990). Effectiveness of reforestation in prevention and control of landsliding during cyclonic storms. In 'Proceedings of International Union of Forest Research Organisations 19th World Congress. Montreal, Canada', 5-11 August 1990. Volume 1 pp. 340-350.
- Power IL, Dodd MB, Thorrold BS (1999). A comparison of pasture and soil moisture under *Acacia melanoxylon* and *Eucalyptus nitens*. *Proceedings of the New Zealand Grassland Association* **61**, 203-207.
- Power IL, Dodd MB, Thorrold BS (2001). Deciduous or evergreen: Does it make a difference to understorey pasture yield and riparian zone management? *Proceedings of the New Zealand Grassland Association* **63**, 121-125.
- Quinn JM, Dodd MB, Thorrold BS (2007). Whatawhata Catchment Management Project: the story so far. *Proceedings of the New Zealand Grassland Association* **69**, 229-233.
- Raymond K (2012). Increase profitability of harvesting small-scale forests on steep terrain. *New Zealand Tree Grower* **33**, 6-9.
- Robertson K, Loza-Balbuena I, Ford-Robertson J (2004). Monitoring and economic factors affecting the economic viability of afforestation for carbon sequestration projects. *Environmental Science and Policy* **7**, 465-475. doi: <http://dx.doi.org/10.1016/j.envsci.2004.07.003>
- Ross DJ, Tate KR, Scott NA, Feltham CW (1999). Land-use change: Effects on soil carbon, nitrogen and phosphorus pools and fluxes in three adjacent ecosystems. *Soil Biology and Biochemistry* **31**, 803-813.
- Ross DJ, Tate KR, Scott NA, Wilde RH, Rodda NJ, Townsend JA (2002). Afforestation of pastures with *Pinus radiata* influences soil carbon and nitrogen pools and mineralisation and microbial properties. *Soil Research* **40**, 1303-1318.
- Russell JR, Betteridge K, Costall DA, Mackay AD (2001). Cattle treading effects on sediment loss and water infiltration. *Journal of Range Management* **54**, 184-190.
- Schon NL, Mackay AD, Gray RAJ, Minor MA (2011). Influence of phosphorus inputs and sheep treading on soil macrofauna and mesofauna in hill pastures. *New Zealand Journal of Agricultural Research* **54**, 83-96. doi: [10.1080/00288233.2011.558904](http://dx.doi.org/10.1080/00288233.2011.558904)
- Scott D, Keogh JM, Cossens GG, Maunsell LA, Floate MJS, Wills BJ, Douglas GB (1985). Limitations to pasture production and choice of species. In 'Using Herbage Cultivars.' (Eds RE Burgess, JL Brock.) pp. 9-15. (Grassland Research and Practice Series No. 3, New Zealand Grassland

- Association: Palmerston North)
- Sheppard JS, Bulloch BT (1986). Management and uses of *Acacia* spp. (wattles) and *Albizia* spp. (brush wattles). In 'Plant materials handbook for soil conservation. Introduced plants.' (Eds CWS Van Kraayenoord, RL Hathaway.) Vol. 2 pp. 7-19. (Water and Soil Miscellaneous Publication No. 94. Water and Soil Directorate, Ministry of Works and Development: Wellington)
- Singh BK, Tate KR, Kolipaka G, Hedley CB, Macdonald CA, Millard P, Murrell JC (2007). Effect of afforestation and reforestation of pastures on the activity and population dynamics of methanotrophic bacteria. *Applied and Environmental Microbiology* **73**, 5153-5161.
- Spiers JJK (1987). 'Logging operations guidelines.' (New Zealand Logging Industry Research Organisation: Rotorua)
- Spinelli R, Magagnotti N, Nati C (2011). Work quality and veneer value recovery of mechanised and manual log-making in Italian poplar plantations. *European Journal of Forest Research* **130**, 737-744.
- Thompson RC, Luckman PG (1993). Performance of biological erosion control in New Zealand soft rock hill terrain. *Agroforestry Systems* **21**, 191-211.
- Turley DB, Chaudhry Q, Watkins RW, Clark JH, Deswarte FEI (2006). Chemical products from temperate forest tree species—Developing strategies for exploitation. *Industrial Crops and Products* **24**, 238-243.
- Vaughan LW (1984). 'Logging and the environment: a review of research findings and management practices.' (New Zealand Logging Industry Research Organisation: Rotorua)
- Wall AJ (2006). The effect of poplar stand density on hill country pastures. PhD thesis, Massey University.
- Wall AJ, Kemp PD, MacKay AD (2006). Predicting pasture production under poplars using canopy closure images. *Proceedings of the New Zealand Grassland Association* **68**, 325-330.
- Wall AJ, Kemp PD, Mackay AD, Power IL (2010). Evaluation of easily measured stand inventory parameters as predictors of PAR transmittance for use in poplar silvopastoral management. *Agriculture, Ecosystems & Environment* **139**, 665-674. doi: <http://dx.doi.org/10.1016/j.agee.2010.10.012>
- White JGH (1999). The farmlands of New Zealand. In 'New Zealand Pasture and Crop Science.' (Eds J White, J Hodgson.) pp. 1-10. (Oxford University Press: Auckland)
- Wilkinson AG (1999). Poplars and willows for soil erosion control in New Zealand. *Biomass and Bioenergy* **16**, 263-274. doi: [10.1016/s0961-9534\(99\)00007-0](https://doi.org/10.1016/s0961-9534(99)00007-0)
- Woollons RC, Manley BR (2012). Examining growth dynamics of *Pinus radiata* plantations at old ages in New Zealand. *Forestry* **85**, 79-86. doi: [10.1093/forestry/cpr059](https://doi.org/10.1093/forestry/cpr059)